

Tutorials for Program Verification
Exercise sheet 1

Definition (poset) Let L be a set and \leq be a binary relation over L . We call (L, \leq) a *poset* if

- \leq is reflexive (i.e., for all $x \in L : x \leq x$),
- \leq is antisymmetric (i.e., for all $x, y \in L : \text{if } x \leq y \text{ and } y \leq x \text{ then } x = y$), and
- \leq is transitive (i.e., for all $x, y, z \in L : \text{if } x \leq y \text{ and } y \leq z \text{ then } x \leq z$).

Definition (galois connection) Let (L_1, \leq_1) and (L_2, \leq_2) be posets, let α be a function from L_1 to L_2 , let γ be a function from L_2 to L_1 . We call the pair (α, γ) a *galois connection* if

$$\text{for all } x \in L_1 \quad \text{for all } y \in L_2 \quad \alpha(x) \leq_2 y \Leftrightarrow x \leq_1 \gamma(y)$$

holds.

Exercise 1: Galois Connection - Examples

2 bonus points

Consider the following table. Each of the seven rows contains an example for posets (L_1, \leq_1) and (L_2, \leq_2) and functions α and γ . In four rows the pair (α, γ) is not a galois connection. Find these four rows and show for each of these four rows that (α, γ) is not a galois connection.

	(L_1, \leq_1)	(L_2, \leq_2)	$\alpha : L_1 \rightarrow L_2$	$\gamma : L_2 \rightarrow L_1$
1)	$(\mathbb{R}, \leq_{\mathbb{R}})$	$(\mathbb{R}, \leq_{\mathbb{R}})$	$\alpha(x) = \lceil x \rceil$	$\gamma(y) = \lfloor y \rfloor$
2)	$(\mathbb{R}, \leq_{\mathbb{R}})$	$(\mathbb{R}, \leq_{\mathbb{R}})$	$\alpha(x) = \lfloor x \rfloor$	$\gamma(y) = \lceil y \rceil$
3)	$(2^{\mathbb{R}}, \subseteq)$	$(2^{\mathbb{R}}, \subseteq)$	$\alpha(X) = \{\lfloor x \rfloor \mid x \in X\}$	$\gamma(Y) = \{\lceil y \rceil \mid y \in Y\}$
4)	$(2^{\mathbb{R}}, \subseteq)$	$(2^{\mathbb{R}_0^+}, \subseteq)$	$\alpha(X) = \{ x \mid x \in X\}$	$\gamma(Y) = \{-y, y \mid y \in Y\}$
5)	$(2^{\mathbb{R}}, \subseteq)$	$(2^{\mathbb{R}}, \subseteq)$	$\alpha(X) = \{ x \mid x \in X\}$	$\gamma(Y) = \{-y, y \mid y \in Y\}$
6)	$(2^{\mathbb{N}}, \subseteq)$	$(2^{\mathbb{N}}, \subseteq)$	$\alpha(X) = \emptyset$	$\gamma(Y) = \mathbb{N}$
7)	$(2^{\mathbb{N}}, \subseteq)$	$(2^{\mathbb{N}}, \subseteq)$	$\alpha(X) = \mathbb{N}$	$\gamma(Y) = \emptyset$

The usual order on real numbers is denoted by $\leq_{\mathbb{R}}$. The unary operators $\lceil \cdot \rceil$, $\lfloor \cdot \rfloor$, and $|\cdot|$ denote the functions for rounding up, rounding down, and absolute value of a real number. The set of all positive real numbers and zero is denoted by \mathbb{R}_0^+ . For a set S the powerset of S is denoted by 2^S .

Exercise 2: Galois Connection - Formalization

2 bonus points

The section Intuition in the Wikipedia article

http://en.wikipedia.org/wiki/Abstract_interpretation

(version from Thu Oct 27, 6pm) informally discusses two abstraction functions on the domain of sets of persons (the sets are ordered by inclusion). Formalize the abstraction, i.e., define the two corresponding abstract domains and the abstraction function α and the concretization function γ . You can assume that we have functions such as:

$$\begin{aligned}SSN &: PERSONS \rightarrow \mathbb{N} \\age &: PERSONS \rightarrow \mathbb{N} \\name &: PERSONS \rightarrow String\end{aligned}$$

Definition (monotonicity) Let (L_1, \leq_1) and (L_2, \leq_2) be posets, we call a function $f : L_1 \rightarrow L_2$ *monotone* if for all $x, x' \in L_1$ $x \leq_1 x'$ implies $f(x) \leq_2 f(x')$.

Exercise 3: Galois Connection - Properties

4+2 bonus points

Let (L_1, \leq_1) and (L_2, \leq_2) be posets and $\alpha : L_1 \rightarrow L_2, \gamma : L_2 \rightarrow L_1$ be functions.

- (a) Show that (α, γ) is a galois connection if and only if
- $\gamma \circ \alpha$ is extensive (i.e., for all $x \in L_1$ $x \leq_1 \gamma(\alpha(x))$),
 - $\alpha \circ \gamma$ is reductive (i.e., for all $y \in L_2$ $\alpha(\gamma(y)) \leq_2 y$),
 - α is monotone,
 - and γ is monotone.
- (b) Let (α, γ) be a galois connection. Show that α is surjective if and only if γ is injective.